

UNIVERSITÄTSBIBLIOTHEK

edoc

Institutional Repository of the University of Basel

University Library

Schoenbeinstrasse 18-20

CH-4056 Basel, Switzerland

<http://edoc.unibas.ch/>

UB | Für alle offen

Year: 2009

The biology of PGC-1 α and its therapeutic potential

Handschin, C.

Posted at edoc, University of Basel

Official URL: <http://edoc.unibas.ch/dok/A5258706>

Originally published as:

Handschin, C.. (2009) *The biology of PGC-1 α and its therapeutic potential*. Trends in pharmacological sciences, Vol. 30, H. 6. S. 322-329.

The biology of PGC-1 α and its therapeutic potential

Christoph Handschin^{1,2}

¹Biozentrum, University of Basel, Klingelbergstrasse 50-70, CH-4056 Basel, Switzerland

²Institute of Physiology and Zurich Center for Integrative Human Physiology (ZIHP), University of Zurich, Winterthurerstrasse 190, CH-8057 Zurich, Switzerland

Published in Trends Pharmacol Sci. 2009 Jun;30(6):322-9. PMID: 19446346doi: 10.1016/j.tips.2009.03.006

Copyright © Elsevier; Trends in Pharmacological Sciences

1 **The biology of PGC-1 α and its therapeutic potential**

2

3 Christoph Handschin^{1,2}

4

5 ¹Biozentrum, University of Basel, Klingelbergstrasse 50-70, CH-4056 Basel,
6 Switzerland

7 ²Institute of Physiology and Zurich Center for Integrative Human Physiology
8 (ZIHP), University of Zurich, Winterthurerstrasse 190, CH-8057 Zurich,
9 Switzerland

10

11 Corresponding author: Handschin, C. (christoph.handschin@unibas.ch)

1 **Abstract**

2 In eukaryotes, cellular and systemic metabolism is primarily controlled by
3 mitochondrial activity. The peroxisome proliferator-activated receptor γ
4 coactivator 1 α (PGC-1 α) is a major regulator of mitochondrial biogenesis and
5 function. Furthermore, PGC-1 α controls many of the phenotypic adaptations of
6 oxidative tissues to external and internal perturbations. In contrast, dysregulated
7 metabolic plasticity is involved in the etiology of numerous diseases. Accordingly,
8 modulation of PGC-1 α levels and activity has recently been proposed as a
9 therapeutic option for several pathologies. However, pharmacological
10 interventions aimed at PGC-1 α have to overcome inherent limitations of targeting
11 a coactivator protein. This review focuses on the recent breakthroughs in the
12 identification of physiological and pathophysiological contexts involving PGC-1 α .
13 In addition, perspectives regarding the therapeutic importance of PGC-1 α -
14 controlled cellular and systemic metabolism are outlined.

15

1 **Introduction**

2

3 Transcriptional changes in gene expression underlie the coordinated regulation
4 of biological programs. These changes are initiated and maintained by the
5 binding of regulatory protein complexes to DNA elements in the enhancer and
6 promoter regions of target genes. Traditionally, DNA-binding transcription factors
7 were thought to be the main regulators of gene expression. However, in recent
8 years, the importance of transcriptional coregulators in the coordination of the
9 expression of genetic programs has been appreciated [1, 2]. The family of
10 peroxisome proliferator-activated receptor γ coactivator-1 (PGC-1) genes
11 illustrates how coactivators respond to environmental cues and subsequently
12 regulate biological processes in a tissue-specific and highly coordinated manner
13 [3-5]. All three members of this small gene family, PGC-1 α , PGC-1 β and PGC-
14 related coactivator (PRC) are strong promoters of mitochondrial biogenesis and
15 oxidative metabolism [6-8]. In contrast to PRC that is ubiquitously expressed,
16 PGC-1 α and PGC-1 β are primarily found in oxidative tissues, including brain,
17 heart, kidney, muscle, liver, brown adipose tissue (BAT) and pancreas. In these
18 organs, PGC-1 α and PGC-1 β have overlapping as well as clearly distinct
19 functions [9]. Moreover, these two coactivators are differently regulated in
20 development and in response to nutritional and other challenges [8, 10].
21 However, whereas our understanding of the physiological role of PGC-1 β
22 remains rudimentary, significant progress in the study of PGC-1 α has been made
23 since its discovery more than a decade ago [11]. This review aims at

1 summarizing recent findings and at highlighting the potential therapeutic
2 applications of PGC-1 α .

4 **PGC-1 α regulates tissue-specific gene expression in health and disease**

6 The PGC-1 α protein is very versatile and coactivates many transcription factors
7 [3-5]. Binding to different partners enables PGC-1 α to regulate distinct biological
8 programs. For example, a combination of coactivation events among PGC-1 α ,
9 the hepatic nuclear factor 4 α (HNF4 α) and the forkhead transcription factor
10 Foxo1 determines the rate of fasting-induced hepatic gluconeogenesis [12, 13].
11 In skeletal muscle, myofibrillar genes are controlled by PGC-1 α -mediated
12 coactivation of myocyte enhancer factor 2 (MEF2) proteins [14]. In most tissues,
13 abnormal regulation of PGC-1 α expression and protein activity results in
14 pathological consequences. Furthermore, single nucleotide polymorphisms
15 (SNP) of *PPARGC1A*, the PGC-1 α gene, have been associated with a diverse
16 set of human diseases (<http://www.geneticassociationdb.nih.gov>). PGC-1 α SNPs
17 are found throughout the protein coding region as well as in the promoter, introns
18 and the 3' untranslated region (3' UTR). The functional consequences of the
19 SNPs are not clearly understood and disease-association of individual SNPs
20 seems highly population- and context-dependent.

22 **Adipose tissue**

1 In BAT, PGC-1 α expression is regulated by cold exposure and β -adrenoreceptor
2 agonists [11]. Subsequently, PGC-1 α coactivates PPAR γ and the thyroid
3 hormone receptor (TR) on the uncoupling protein 1 (UCP-1), type 2 deiodinase
4 (DIO2) and other BAT-specific gene promoters [11]. BAT vs. white adipose tissue
5 (WAT) selectivity of gene expression is controlled by the distinct assembly of
6 PGC-1 α / PR(PRD1-BF1-RIZ1 homologous)-domain-containing protein (Prdm16)
7 / C-terminal binding protein-1 (CtBP1) protein complexes in these two tissues
8 [15]. PGC-1 α function is essential for BAT-mediated adaptive thermogenesis *in*
9 *vitro* and *in vivo* [9, 16, 17]. Thus, when exposed to cold, PGC-1 α knockout mice
10 rapidly become hypothermic and die after prolonged exposure [16, 17].

11
12 Levels of PGC-1 α in WAT are much lower than those in BAT and the
13 physiological role of PGC-1 α in this tissue is unclear [11]. PGC-1 α expression is
14 further reduced in the subcutaneous fat of morbidly obese and of insulin resistant
15 patients, respectively [18, 19]. In contrast, thiazolidinedione treatment and
16 hyperleptinemia elevate PGC-1 α transcription in WAT; this induction mediates
17 some of the beneficial effects of this class of anti-diabetic drugs and of leptin,
18 respectively [20, 21]. Ectopic expression of PGC-1 α in a WAT context results in
19 the induction of BAT-specific genes such as UCP-1 and an increase in
20 mitochondrial activity [11, 22].

21
22 *Therapeutic potential:* Augmenting the function of the remaining BAT in adult
23 humans or pushing WAT towards BAT-specific gene expression by elevating

PGC-1 α could tilt energy balance towards expenditure and thereby reduce obesity (Fig. 1). Proof-of-concept for a therapeutic effect of augmented BAT content is provided by chronic β 3-adrenoreceptor stimulation resulting in a lean and healthy phenotype in animal models [23]. However, it remains to be tested whether the limited capacity of PGC-1 α to increase BAT gene expression in WAT is sufficient or if other factors such as Prdm16 are required to achieve a therapeutic effect [24].

Liver

The first peak of PGC-1 α gene expression is observed in the liver at birth [10]. In the adult, fasting and glucagon are the main drivers of hepatic PGC-1 α transcription [12, 25]. PGC-1 α regulates most of the metabolic changes that occur during the transition of a fed to a fasted liver, including gluconeogenesis, fatty acid β -oxidation, ketogenesis and heme biosynthesis [12, 13, 26, 27]. Studies with knockout animals and adenovirally delivered shRNA constructs revealed a blunted hepatic fasting response in the absence of adequate levels of PGC-1 α [17, 27, 28]. Accordingly, these mice develop a fasting hypoglycemia and hepatic steatosis [17, 28].

Therapeutic potential: An increase in PGC-1 α levels in the liver might obviously alleviate hypoglycemia and hepatic steatosis. However, in contrast to the induction sought after in most other tissues, a repression of PGC-1 α activity

holds more promise for therapy in the liver. First, hepatic PGC-1 α expression is elevated in mouse models for type 1 and type 2 diabetes and probably contributes to the unchecked glucose production and hyperglycemia in diabetes [12]. Normalization of PGC-1 α levels might thus help to control hepatic gluconeogenesis in these patients. Individuals with acute hepatic porphyrias, diseases caused by mutations in the heme biosynthetic pathway, often suffer from fasting-induced acute attacks that can be alleviated by glucose and heme administration [29]. PGC-1 α has been identified as one of the main culprits in the precipitation of acute porphyric attacks in fasting [27]. These attacks might be prevented by pharmacological interventions that thwart the induction or activity of PGC-1 α [29, 30]. Furthermore, PGC-1 α regulates the genes encoding homocysteine synthesis enzymes in the fasted liver [31]. High plasma homocysteine is an independent risk factor for the development of cardiovascular diseases. Accordingly, inhibition of PGC-1 α could result in a drop of plasma homocysteine levels and thereby a reduction of the risk for cardiovascular diseases [31]. Finally, the hepatitis B virus (HBV) uses the transcriptional machinery that regulates the hepatic fasting response for amplification [32]. Accordingly, HBV gene expression and life cycle, and thus the viral-host interaction, are under the control of PGC-1 α [32]. Repression of PGC-1 α activity might present a new anti-viral therapy for hepatitis B and help to restrict the viral load in this disease.

Brain

1

2 In the developing brain, PGC-1 α transcription peaks two weeks postnatally in
3 many regions, a period of significant metabolic changes, mitochondrial
4 biogenesis and synaptic remodeling [33]. The highest PGC-1 α concentrations
5 are found in γ -aminobutyric acid (GABA)-positive neurons in the cortex,
6 hippocampus and cerebellum [33]. One of the existing global PGC-1 α knockout
7 line exhibits increased anxiety [17], whereas a second model shows a profound
8 hyperactivity [16]. Other behavioral changes include hind limb claspings, dystonic
9 posturing, an exaggerated startle response and stimulus-induced myoclonus
10 [16]. Besides the behavioral abnormalities, spongiform-like vacuolization events
11 are observed in the dorsolateral striatum [16], the vicinity of pyramidal neurons
12 [17], and more sporadically, in other regions of the brain of PGC-1 α knockout
13 mice [16]. Furthermore, the substantia nigra and the CA1 neurons of the
14 hippocampus are more susceptible to degenerative events triggered by the
15 reactive oxygen species (ROS)-generating 1-methyl-4-phenyl-1,2,3,6-
16 tetrahydropyridine (MPTP) and kainic acid, respectively, in these animals [34].
17 These findings suggest an important involvement of PGC-1 α in neuronal
18 maintenance and function.

19

20 *Therapeutic potential:* Mitochondrial dysfunction and oxidative stress are
21 associated with many neurodegenerative disorders, including Parkinson's,
22 Alzheimer's, Huntington's, Friedreich's ataxia and amyotrophic lateral disorder
23 (ALS) [35, 36]. PGC-1 α coordinately regulates the entire mitochondrial program

1 concomitant with an increase in the ROS detoxification system [34]. Reduced
2 PGC-1 α expression in Huntington's patients, the association of PGC-1 α SNPs
3 with the age of onset of Huntington's [37] and the Huntington's-like phenotype in
4 PGC-1 α knockout suggest that PGC-1 α is crucial for the maintenance of proper
5 striatal function [16, 38, 39]. A normalization of PGC-1 α expression in the
6 striatum of Huntington's patients might thus constitute a promising therapeutic
7 option [40-42]. In addition, the wide-spread neuronal lesions and the
8 susceptibility to chemically-induced neurodegeneration in different regions of the
9 brain of PGC-1 α knockout animals imply that PGC-1 α has a broader role in
10 neuroprotection beyond that related to Huntington's disease [16, 34].

12 **Skeletal muscle**

14 In the contracting muscle fiber, the main signaling pathways converge on PGC-
15 1 α to increase expression levels and the activity of this coactivator [3, 43, 44].
16 For example, p38 mitogen-activated protein kinase (p38 MAPK) and AMP-
17 dependent kinase (AMPK) are rapidly activated in exercise and subsequently
18 phosphorylate the PGC-1 α protein. In addition, PGC-1 α transcription is regulated
19 by the motor neuron-induced rise in intracellular calcium, AMPK, β 2-
20 adrenoreceptor signaling, nitric oxide and thyroid hormone [5]. As a
21 consequence, PGC-1 α levels in skeletal muscle are elevated following bouts of
22 endurance exercise. PGC-1 α is primarily found in type I and IIa slow-twitch, high
23 endurance muscle fibers and regulates the adaptations to exercise [14]. In fact,

1 ectopic expression of PGC-1 α in skeletal muscle is sufficient to promote a fiber-
2 type switching towards oxidative muscle fibers and bestow a trained phenotype
3 onto mice [14, 45].

4
5 Abnormally low PGC-1 α levels have been described in skeletal muscle of type 2
6 diabetic patients and physically inactive individuals [46, 47]. Mice with reduced or
7 ablated PGC-1 α gene expression suffer from a decreased exercise capacity,
8 abnormal systemic glucose and insulin homeostases, systemic inflammation and
9 activity-dependent fiber damage [48, 49]. Paradoxically, transgenic mice with
10 elevated PGC-1 α levels in muscle are prone to develop peripheral insulin
11 resistance on a high fat-containing diet [50].

12
13 *Therapeutic potential:* Many diseases are associated with inactive skeletal
14 muscle [51]. In animal models, transgenic elevation of PGC-1 α in muscle blunts
15 inactivity-induced fiber atrophy [52], reduces muscle wasting triggered by the
16 statin class of drugs [53], ameliorates Duchenne muscular dystrophy [54] and a
17 specific form of mitochondrial myopathy [55]. Since PGC-1 α promotes an
18 exercised phenotype in muscle, the therapeutic potential of PGC-1 α against
19 diseases that are associated with dysregulated muscle function is extremely
20 broad.

21
22 PGC-1 α controls the transcription of several genes encoding oxidative
23 phosphorylation (OXPHOS) genes that are dysregulated in muscle of type 2

1 diabetic patients [46, 47]. The specific disruption of the interaction between PGC-
2 1α and the estrogen-related receptor α (ERR α) using an inverse agonist in
3 muscle cells in culture promotes a change in gene expression similar to that
4 observed in muscle of diabetic patients [56]. Activation of the PGC- 1α /ERR α axis
5 could thus be a novel therapeutic approach against insulin resistance and type 2
6 diabetes [57]. However, the data from gain- and loss-of-function animal models of
7 PGC- 1α in muscle imply a complicated relationship between the expression of
8 this coactivator and systemic glucose and insulin levels that has yet to be
9 clarified [49, 50].

11 **Heart**

13 In the heart, PGC- 1α transcription is induced at birth and correlates with
14 metabolic maturation and remodeling [58]. As in other tissues, cardiac PGC- 1α
15 strongly promotes mitochondrial function and fatty acid β -oxidation [59]. In
16 several mouse models for heart disease accompanied with a substrate switch
17 from fatty acid to glucose utilization, expression of PGC- 1α is reduced [60]. At
18 baseline, heart function of PGC- 1α knockout mice appears normal [17, 61].
19 However, the cardiac reserve under stress conditions is impaired in mice with an
20 ablated PGC- 1α gene [61] and accordingly, these mice have a diminished
21 cardiac capacity in exercise [17]. In addition, prolonged pressure overload by
22 transverse aortic constriction precipitates ventricular dysfunction and clinical
23 signs of heart failure in PGC- 1α knockout animals [62]. Interestingly, ectopic

1 expression of PGC-1 α in the heart at superphysiological levels also results in
2 cardiomyopathy and heart failure in mice [58].

3
4 *Therapeutic potential:* Moderate elevation or normalization of PGC-1 α expression
5 in the failing heart might be sufficient to switch substrate usage from glucose
6 back to fatty acids and restore adequate energy production [62]. Thereby, PGC-
7 1 α could be cardioprotective against certain insults [60]. In contrast, insulin-
8 resistant hearts rely heavily on fatty acid oxidation and there, PGC-1 α levels are
9 elevated [63]. It is unclear if upregulation of cardiac PGC-1 α in insulin resistance
10 is an adaptive or a maladaptive process. In the former case, a reduction of PGC-
11 1 α expression should be the therapeutic goal. However, if the increase of PGC-
12 1 α is an adaptive process to cope with the excess amount of fatty acids, a further
13 elevation might be therapeutically beneficial [3, 60].

14 15 **Pancreas**

16
17 In obese mice, insulin resistant animal models and in a partial pancreatectomy as
18 a model for β cell decompensation, the transcriptional rate of the PGC-1 α gene in
19 the pancreas is increased over the normally low basal levels [64]. In pancreatic
20 islets, PGC-1 α prevents membrane polarization and induces glucose-6-
21 phosphatase and thereby reduces insulin secretion [64]. Notably, a second study
22 of PGC-1 α in human pancreatic islets resulted in opposite findings [65]. A
23 common PGC-1 α SNP was associated with reduced PGC-1 α expression and

1 insulin secretion in islets of tissue donors [65]. In addition, silencing of PGC-1 α
2 gene expression in isolated human islets likewise led to impaired insulin
3 secretion [65].

4
5 *Therapeutic potential:* At the moment, the diametrically opposed results of these
6 two studies are difficult to accommodate. Until these issues are resolved, it is
7 unclear whether a suppression or an induction of PGC-1 α activity in pancreatic
8 islets is the more promising avenue to restore insulin secretion in type 1 and type
9 2 diabetes.

10 11 **Bone and cartilage**

12
13 Parathyroid hormone-induced cAMP signaling is a major pathway in osteoclast
14 activation [66]. PGC-1 α is a primary target gene of this signaling cascade and
15 synergistically with Nurr1, a cAMP-induced orphan nuclear receptor, increases
16 the transcription of osteopontin and osteocalcin, two key genes in bone formation
17 [66]. Given the significant role of ERR α , one of the strongest interaction partners
18 of PGC-1 α , in osteoblasts and osteoclasts [67] and the coactivation of the
19 vitamin D receptor by PGC-1 α [68], an important function for PGC-1 α in
20 osteogenesis seems likely. These findings are intriguing in regard to a report
21 showing a role of PGC-1 α in chondrogenesis [69] suggesting an even broader
22 implication for PGC-1 α in the developing skeleton.

1 *Therapeutic options:* An activation of PGC-1 α might be useful in osteoporosis or
2 other disease associated with reduced bone density and morphology [66].
3 Similarly, PGC-1 α could be a promising target for molecular engineering of
4 cartilage in the clinical setting, e.g., osteoarthritis and other chondrodystrophies
5 [69].

6

7 **Problems, pitfalls and opportunities**

8

9 As a transcriptional coactivator, PGC-1 α lacks functional DNA- and ligand-
10 binding domains and therefore is not amenable to direct pharmacological
11 intervention. Strategies to alter the availability of PGC-1 α thus have to aim at
12 transcriptional regulation of the gene, modifications of the protein or the
13 interaction with binding partners. Modulation of PGC-1 α transcription is
14 hampered by the tight regulation of cellular metabolism. Accordingly, despite
15 different screening efforts [70, 71], clinically interesting compounds that
16 persistently induce PGC-1 α transcription are elusive. Optimally, such drugs
17 would act in a tissue-specific manner to circumvent unwanted side-effects by
18 elevating PGC-1 α in non-target tissues. For example, an elevation of PGC-1 α in
19 WAT by thiazolidinedione drugs has been reported [20]; however, it is unclear
20 whether this induction is direct, secondary to other events or even sufficient to
21 contribute to the therapeutic effect of this class of drugs.

22

1 Posttranslational modifications of the PGC-1 α protein alter the half-life and the
2 specificity towards binding partners (Fig. 2). For example, PGC-1 α is
3 phosphorylated by AMPK at two different sites [44]. Metformin, a clinically used
4 anti-diabetic drug mediates at least some of its therapeutic effect through
5 activation of AMPK. Furthermore, deacetylation of 13 lysine residues on PGC-1 α
6 by the mammalian silencing information regulator 2-ortholog SIRT1 increases
7 PGC-1 α activity and thereby targets gene expression in muscle and liver [72, 73].
8 Resveratrol and other, more specific SIRT1 activators are promising drugs
9 against aging and aging-related diseases acting partially through PGC-1 α [74].
10 Other known modifications of PGC-1 α include phosphorylations by p38 MAPK
11 [75, 76] and Akt2/protein kinase B [77], methylation by protein arginine
12 methyltransferase 1 (PRMT1) [78], ubiquitination [79] and O-linked β -N-
13 acetylglucosamination [80]. Proteins with such enzymatic activities are attractive
14 drug targets for which specific activators and inhibitors can be designed.
15 Following the posttranslational modifications that modify the specificity of PGC-
16 1 α , the transcription of selective groups of target genes is altered (Fig. 2). Thus,
17 pharmacological manipulation of these enzymes upstream of PGC-1 α might
18 result in a fine-tuned modulation of PGC-1 α activity in a tissue- and target gene-
19 specific manner [5].

20
21 PGC-1 α regulates biological programs by interacting with different transcriptional
22 complexes. For instance, the strong potentiation of the transcriptional activity of
23 ERR α by PGC-1 α is responsible for mitochondrial OXPHOS gene expression in

1 muscle and other tissues and a specific disruption of this interaction selectively
2 reduces the expression of these, but not other PGC-1 α target genes [56]. The
3 use of partial agonists for ligand-binding transcription factors that modulate the
4 binding of individual coactivators [81] or of compounds that selectively disrupt or
5 enhance protein-protein interactions [56] could thus be another approach for a
6 specific modulation of PGC-1 α activity. Furthermore, pharmacological
7 interventions targeting the assembly of the PGC-1 α -containing transcriptional
8 complex that includes the p160 myb-binding protein [82], histone acetyl-
9 transferase enzymes (HAT) [83], members of the TRAP/DRIP/mediator complex
10 [84] and other proteins [85] are alternative strategies with therapeutic potential.

11
12 Finally, indiscriminate changes in the level of PGC-1 α in either direction are most
13 likely equally detrimental (Fig. 3). For example, cardiomyopathy and heart failure
14 develop with sub- and superphysiological levels of PGC-1 α [58, 62]. In skeletal
15 muscle, an increased exercise capacity was observed in a mouse line with
16 moderate transgenic expression of PGC-1 α [14]. However, the roughly ten-fold
17 increase in PGC-1 α transcript levels in these animals might already be excessive
18 for normal insulin sensitivity [50]. In contrast, an even more modest elevation of
19 PGC-1 α as demonstrated in an *in vivo* transfection experiment clearly increased
20 insulin sensitivity in skeletal muscle [86]. Higher expressing mouse lines suffer
21 from displacement of the contractile apparatus by excess mitochondria, weight
22 loss, fiber atrophy and muscle wasting [14, 87], reminiscent of the fiber damage
23 of muscle-specific knockout animals [48, 49]. Modulation of PGC-1 α thus has to

aim at either a moderate alteration or a normalization of PGC-1 α activity within a therapeutically beneficial window that might be tissue- and context-dependent.

Perspectives

The transcriptional coactivator PGC-1 α plays a key role in maintaining cellular metabolism. Dysregulation of gene expression and gene polymorphisms of PGC-1 α have accordingly been found in a wide variety of different pathological contexts. Furthermore, therapeutic efficacy of PGC-1 α modulation has been demonstrated in animal models for different diseases. Thus, pharmacological regulation of PGC-1 α expression and activity might be a promising novel approach for the prevention and therapy of a number of pathologies despite the inherent difficulties of targeting a coactivator. On the other hand, coactivators exhibit some features that could be exploited for clinical use. By coordinating different steps in biological programs through the integration of the activity of many transcription factors, coactivators have kinetic advantages in controlling biological programs over transcription factor-mediated regulation [1]. Furthermore, combinatorial combinations of posttranslational modifications of coregulators exponentially increase the degree of specification of these proteins [2, 5]. Thus, a selective targeting of PGC-1 α could potentially result in a fine-tuned and highly specific response, which is unattainable by modulation of transcription factors. Finally, modulation of PGC-1 α in one tissue might trigger distal effects in other organs. For example, observations from muscle-specific

1 loss-of-function mouse lines suggest that PGC-1 α links muscle function to
2 systemic inflammation and ultimately, the risk of developing many chronic
3 diseases [48, 49, 51]. If these findings are substantiated, then modulation of
4 PGC-1 α might have a much broader therapeutic applicability [51]. Obviously, big
5 gaps of knowledge about the function of PGC-1 α , and even more about PGC-1 β
6 and PRC exist. Hopefully, a more comprehensive understanding of the
7 therapeutic potential of the PGC-1 family of coactivators will emerge from future
8 studies addressing mechanistic and physiological aspects of these intriguing
9 proteins.

1 **Acknowledgments**

2

3 I thank my colleagues for discussions, ideas and suggestions for writing this
4 manuscript and Christian Gasser for help with the artwork. I apologize for the
5 omission of several important contributions due to space constraints. Work in my
6 laboratory related to this manuscript has been supported by the Swiss National
7 Science Foundation (SNF PP00A-110746), the Muscular Dystrophy Association
8 USA (MDA), the SwissLife “Jubiläumsstiftung für Volksgesundheit und
9 medizinische Forschung”, the Swiss Society for Research on Muscle Diseases
10 (SSEM), the Swiss Diabetes Association, the Roche Research Foundation, the
11 Zurich Center for Integrative Human Physiology (ZIHP) and the Universities of
12 Basel and Zurich.

13

References

1. Spiegelman, B.M., and Heinrich, R. (2004) Biological control through regulated transcriptional coactivators. *Cell* 119, 157-167
2. Lonard, D.M., and O'Malley B, W. (2007) Nuclear receptor coregulators: judges, juries, and executioners of cellular regulation. *Mol Cell* 27, 691-700
3. Finck, B.N., and Kelly, D.P. (2006) PGC-1 coactivators: inducible regulators of energy metabolism in health and disease. *J Clin Invest* 116, 615-622
4. Lin, J., et al. (2005) Metabolic control through the PGC-1 family of transcription coactivators. *Cell Metab* 1, 361-370
5. Handschin, C., and Spiegelman, B.M. (2006) PGC-1 coactivators, energy homeostasis, and metabolism. *Endocr Rev* 27, 728-735
6. Wu, Z., et al. (1999) Mechanisms controlling mitochondrial biogenesis and respiration through the thermogenic coactivator PGC-1. *Cell* 98, 115-124
7. Andersson, U., and Scarpulla, R.C. (2001) Pgc-1-related coactivator, a novel, serum-inducible coactivator of nuclear respiratory factor 1-dependent transcription in mammalian cells. *Mol Cell Biol* 21, 3738-3749
8. Lin, J., et al. (2002) Peroxisome proliferator-activated receptor gamma coactivator 1beta (PGC-1beta), a novel PGC-1-related transcription coactivator associated with host cell factor. *J Biol Chem* 277, 1645-1648
9. Uldry, M., et al. (2006) Complementary action of the PGC-1 coactivators in mitochondrial biogenesis and brown fat differentiation. *Cell Metab* 3, 333-341

- 1 10. Lin, J., *et al.* (2003) PGC-1beta in the regulation of hepatic glucose and
2 energy metabolism. *J Biol Chem* 278, 30843-30848
- 3 11. Puigserver, P., *et al.* (1998) A cold-inducible coactivator of nuclear
4 receptors linked to adaptive thermogenesis. *Cell* 92, 829-839
- 5 12. Yoon, J.C., *et al.* (2001) Control of hepatic gluconeogenesis through the
6 transcriptional coactivator PGC-1. *Nature* 413, 131-138
- 7 13. Puigserver, P., *et al.* (2003) Insulin-regulated hepatic gluconeogenesis
8 through FOXO1-PGC-1alpha interaction. *Nature* 423, 550-555
- 9 14. Lin, J., *et al.* (2002) Transcriptional co-activator PGC-1 alpha drives the
10 formation of slow-twitch muscle fibres. *Nature* 418, 797-801
- 11 15. Kajimura, S., *et al.* (2008) Regulation of the brown and white fat gene
12 programs through a PRDM16/CtBP transcriptional complex. *Genes Dev* 22,
13 1397-1409
- 14 16. Lin, J., *et al.* (2004) Defects in adaptive energy metabolism with CNS-
15 linked hyperactivity in PGC-1alpha null mice. *Cell* 119, 121-135
- 16 17. Leone, T.C., *et al.* (2005) PGC-1alpha deficiency causes multi-system
17 energy metabolic derangements: muscle dysfunction, abnormal weight control
18 and hepatic steatosis. *PLoS Biol* 3, e101
- 19 18. Semple, R.K., *et al.* (2004) Expression of the thermogenic nuclear
20 hormone receptor coactivator PGC-1alpha is reduced in the adipose tissue of
21 morbidly obese subjects. *Int J Obes Relat Metab Disord* 28, 176-179

- 1 19. Yang, X., *et al.* (2003) Reduced expression of FOXC2 and brown
2 adipogenic genes in human subjects with insulin resistance. *Obes Res* 11, 1182-
3 1191
- 4 20. Wilson-Fritch, L., *et al.* (2004) Mitochondrial remodeling in adipose tissue
5 associated with obesity and treatment with rosiglitazone. *J Clin Invest* 114, 1281-
6 1289
- 7 21. Lee, Y., *et al.* (2002) PPAR alpha is necessary for the lipopenic action of
8 hyperleptinemia on white adipose and liver tissue. *Proc Natl Acad Sci U S A* 99,
9 11848-11853
- 10 22. Tiraby, C., *et al.* (2003) Acquirement of brown fat cell features by human
11 white adipocytes. *J Biol Chem* 278, 33370-33376
- 12 23. Ghorbani, M., *et al.* (1997) Hypertrophy of brown adipocytes in brown and
13 white adipose tissues and reversal of diet-induced obesity in rats treated with a
14 beta3-adrenoceptor agonist. *Biochem Pharmacol* 54, 121-131
- 15 24. Seale, P., *et al.* (2007) Transcriptional control of brown fat determination
16 by PRDM16. *Cell Metab* 6, 38-54
- 17 25. Herzig, S., *et al.* (2001) CREB regulates hepatic gluconeogenesis through
18 the coactivator PGC-1. *Nature* 413, 179-183
- 19 26. Rhee, J., *et al.* (2003) Regulation of hepatic fasting response by
20 PPARgamma coactivator-1alpha (PGC-1): Requirement for hepatocyte nuclear
21 factor 4alpha in gluconeogenesis. *Proc Natl Acad Sci U S A* 100, 4012-4017
- 22 27. Handschin, C., *et al.* (2005) Nutritional regulation of hepatic heme
23 biosynthesis and porphyria through PGC-1alpha. *Cell* 122, 505-515

- 1 28. Koo, S.H., *et al.* (2004) PGC-1 promotes insulin resistance in liver through
2 PPAR-alpha-dependent induction of TRB-3. *Nat Med* 10, 530-534
- 3 29. Phillips, J.D., and Kushner, J.P. (2005) Fast track to the porphyrias. *Nat*
4 *Med* 11, 1049-1050
- 5 30. Li, D. (2005) PGC-1alpha: looking behind the sweet treat for porphyria.
6 *Cell* 122, 487-489
- 7 31. Li, S., *et al.* (2009) Regulation of homocysteine homeostasis through the
8 transcriptional coactivator PGC-1{alpha}. *Am J Physiol Endocrinol Metab* 296,
9 E543-548
- 10 32. Shlomai, A., *et al.* (2006) PGC-1alpha controls hepatitis B virus through
11 nutritional signals. *Proc Natl Acad Sci U S A* 103, 16003-16008
- 12 33. Cowell, R.M., *et al.* (2007) Localization of the transcriptional coactivator
13 PGC-1alpha to GABAergic neurons during maturation of the rat brain. *The*
14 *Journal of comparative neurology* 502, 1-18
- 15 34. St-Pierre, J., *et al.* (2006) Suppression of Reactive Oxygen Species and
16 Neurodegeneration by the PGC-1 Transcriptional Coactivators. *Cell* 127, 397-408
- 17 35. Lin, M.T., and Beal, M.F. (2006) Mitochondrial dysfunction and oxidative
18 stress in neurodegenerative diseases. *Nature* 443, 787-795
- 19 36. Chaturvedi, R.K., and Beal, M.F. (2008) Mitochondrial approaches for
20 neuroprotection. *Ann N Y Acad Sci* 1147, 395-412
- 21 37. Taherzadeh-Fard, E., *et al.* (2009) PGC-1alpha as modifier of onset age in
22 Huntington disease. *Molecular neurodegeneration* 4, 10

- 1 38. Cui, L., *et al.* (2006) Transcriptional repression of PGC-1alpha by mutant
2 huntingtin leads to mitochondrial dysfunction and neurodegeneration. *Cell* 127,
3 59-69
- 4 39. Weydt, P., *et al.* (2006) Thermoregulatory and metabolic defects in
5 Huntington's disease transgenic mice implicate PGC-1alpha in Huntington's
6 disease neurodegeneration. *Cell Metab* 4, 349-362
- 7 40. McGill, J.K., and Beal, M.F. (2006) PGC-1alpha, a new therapeutic target
8 in Huntington's disease? *Cell* 127, 465-468
- 9 41. Greenamyre, J.T. (2007) Huntington's disease--making connections. *N*
10 *Engl J Med* 356, 518-520
- 11 42. Ross, C.A., and Thompson, L.M. (2006) Transcription meets metabolism
12 in neurodegeneration. *Nat Med* 12, 1239-1241
- 13 43. Handschin, C., *et al.* (2003) An autoregulatory loop controls peroxisome
14 proliferator-activated receptor gamma coactivator 1alpha expression in muscle.
15 *Proc Natl Acad Sci U S A* 100, 7111-7116
- 16 44. Jager, S., *et al.* (2007) AMP-activated protein kinase (AMPK) action in
17 skeletal muscle via direct phosphorylation of PGC-1alpha. *Proc Natl Acad Sci U*
18 *S A* 104, 12017-12022
- 19 45. Calvo, J.A., *et al.* (2008) Muscle-specific expression of PPAR{gamma}
20 coactivator-1{alpha} improves exercise performance and increases peak oxygen
21 uptake. *J Appl Physiol* 104, 1304-1312

- 1 46. Patti, M.E., *et al.* (2003) Coordinated reduction of genes of oxidative
2 metabolism in humans with insulin resistance and diabetes: Potential role of
3 PGC1 and NRF1. *Proc Natl Acad Sci U S A* 100, 8466-8471
- 4 47. Mootha, V.K., *et al.* (2003) PGC-1alpha-responsive genes involved in
5 oxidative phosphorylation are coordinately downregulated in human diabetes.
6 *Nat Genet* 34, 267-273
- 7 48. Handschin, C., *et al.* (2007) Skeletal muscle fiber-type switching, exercise
8 intolerance, and myopathy in PGC-1alpha muscle-specific knock-out animals. *J*
9 *Biol Chem* 282, 30014-30021
- 10 49. Handschin, C., *et al.* (2007) Abnormal glucose homeostasis in skeletal
11 muscle-specific PGC-1alpha knockout mice reveals skeletal muscle-pancreatic
12 beta cell crosstalk. *J Clin Invest* 117, 3463-3474
- 13 50. Choi, C.S., *et al.* (2008) Paradoxical effects of increased expression of
14 PGC-1alpha on muscle mitochondrial function and insulin-stimulated muscle
15 glucose metabolism. *Proc Natl Acad Sci U S A* 105, 19926-19931
- 16 51. Handschin, C., and Spiegelman, B.M. (2008) The role of exercise and
17 PGC1alpha in inflammation and chronic disease. *Nature* 454, 463-469
- 18 52. Sandri, M., *et al.* (2006) PGC-1{alpha} protects skeletal muscle from
19 atrophy by suppressing FoxO3 action and atrophy-specific gene transcription.
20 *Proc Natl Acad Sci U S A* 103, 16260-16265
- 21 53. Hanai, J.I., *et al.* (2007) The muscle-specific ubiquitin ligase atrogin-
22 1/MAFbx mediates statin-induced muscle toxicity. *J Clin Invest* 117, 3940-3951

- 1 54. Handschin, C., *et al.* (2007) PGC-1alpha regulates the neuromuscular
2 junction program and ameliorates Duchenne muscular dystrophy. *Genes Dev* 21,
3 770-783
- 4 55. Wenz, T., *et al.* (2008) Activation of the PPAR/PGC-1alpha pathway
5 prevents a bioenergetic deficit and effectively improves a mitochondrial myopathy
6 phenotype. *Cell Metab* 8, 249-256
- 7 56. Mootha, V.K., *et al.* (2004) Erralpha and Gabpa/b specify PGC-1alpha-
8 dependent oxidative phosphorylation gene expression that is altered in diabetic
9 muscle. *Proc Natl Acad Sci U S A* 101, 6570-6575
- 10 57. Handschin, C., and Mootha, V.K. (2005) Estrogen-related receptor alpha
11 (ERRalpha): A novel target in type 2 diabetes. *Drug Discov Today Ther Strateg*
12 2, 151-156
- 13 58. Lehman, J.J., *et al.* (2000) Peroxisome proliferator-activated receptor
14 gamma coactivator-1 promotes cardiac mitochondrial biogenesis. *J Clin Invest*
15 106, 847-856
- 16 59. Huss, J.M., and Kelly, D.P. (2004) Nuclear receptor signaling and cardiac
17 energetics. *Circ Res* 95, 568-578
- 18 60. Huss, J.M., and Kelly, D.P. (2005) Mitochondrial energy metabolism in
19 heart failure: a question of balance. *J Clin Invest* 115, 547-555
- 20 61. Arany, Z., *et al.* (2005) Transcriptional coactivator PGC-1 alpha controls
21 the energy state and contractile function of cardiac muscle. *Cell Metab* 1, 259-
22 271

- 1 62. Arany, Z., *et al.* (2006) Transverse aortic constriction leads to accelerated
2 heart failure in mice lacking PPAR-gamma coactivator 1alpha. *Proc Natl Acad*
3 *Sci U S A* 103, 10086-10091
- 4 63. Finck, B.N., *et al.* (2002) The cardiac phenotype induced by PPARalpha
5 overexpression mimics that caused by diabetes mellitus. *J Clin Invest* 109, 121-
6 130
- 7 64. Yoon, J.C., *et al.* (2003) Suppression of beta cell energy metabolism and
8 insulin release by PGC-1alpha. *Dev Cell* 5, 73-83
- 9 65. Ling, C., *et al.* (2008) Epigenetic regulation of PPARGC1A in human type
10 2 diabetic islets and effect on insulin secretion. *Diabetologia* 51, 615-622
- 11 66. Nervina, J.M., *et al.* (2006) PGC-1alpha is induced by parathyroid
12 hormone and coactivates Nurr1-mediated promoter activity in osteoblasts. *Bone*
13 39, 1018-1025
- 14 67. Zirngibl, R.A., *et al.* (2008) Estrogen receptor-related receptor alpha
15 (ERRalpha) regulates osteopontin expression through a non-canonical ERRalpha
16 response element in a cell context-dependent manner. *J Mol Endocrinol* 40, 61-
17 73
- 18 68. Savkur, R.S., *et al.* (2005) Coactivation of the human vitamin D receptor
19 by the peroxisome proliferator-activated receptor gamma coactivator-1 alpha. *Mol*
20 *Pharmacol* 68, 511-517
- 21 69. Kawakami, Y., *et al.* (2005) Transcriptional coactivator PGC-1alpha
22 regulates chondrogenesis via association with Sox9. *Proc Natl Acad Sci U S A*
23 102, 2414-2419

- 1 70. Arany, Z., *et al.* (2008) Gene expression-based screening identifies
2 microtubule inhibitors as inducers of PGC-1alpha and oxidative phosphorylation.
3 *Proc Natl Acad Sci U S A* 105, 4721-4726
- 4 71. Wagner, B.K., *et al.* (2008) Large-scale chemical dissection of
5 mitochondrial function. *Nat Biotechnol* 26, 343-351
- 6 72. Rodgers, J.T., *et al.* (2005) Nutrient control of glucose homeostasis
7 through a complex of PGC-1alpha and SIRT1. *Nature* 434, 113-118
- 8 73. Gerhart-Hines, Z., *et al.* (2007) Metabolic control of muscle mitochondrial
9 function and fatty acid oxidation through SIRT1/PGC-1alpha. *Embo J* 26, 1913-
10 1923
- 11 74. Lagouge, M., *et al.* (2006) Resveratrol improves mitochondrial function
12 and protects against metabolic disease by activating SIRT1 and PGC-1alpha.
13 *Cell* 127, 1109-1122
- 14 75. Puigserver, P., *et al.* (2001) Cytokine stimulation of energy expenditure
15 through p38 MAP kinase activation of PPARgamma coactivator-1. *Mol Cell* 8,
16 971-982
- 17 76. Knutti, D., *et al.* (2001) Regulation of the transcriptional coactivator PGC-1
18 via MAPK-sensitive interaction with a repressor. *Proc Natl Acad Sci U S A* 98,
19 9713-9718
- 20 77. Li, X., *et al.* (2007) Akt/PKB regulates hepatic metabolism by directly
21 inhibiting PGC-1alpha transcription coactivator. *Nature* 447, 1012-1016
- 22 78. Teyssier, C., *et al.* (2005) Activation of nuclear receptor coactivator PGC-
23 1alpha by arginine methylation. *Genes Dev* 19, 1466-1473

- 1 79. Sano, M., *et al.* (2007) Intramolecular control of protein stability,
2 subnuclear compartmentalization, and coactivator function of peroxisome
3 proliferator-activated receptor gamma coactivator 1alpha. *J Biol Chem* 282,
4 25970-25980
- 5 80. Housley, M.P., *et al.* (2009) A PGC-1{alpha}-O-GlcNAc Transferase
6 Complex Regulates FoxO Transcription Factor Activity in Response to Glucose. *J*
7 *Biol Chem* 284, 5148-5157
- 8 81. Coghlan, M.J., *et al.* (2003) A novel antiinflammatory maintains
9 glucocorticoid efficacy with reduced side effects. *Mol Endocrinol* 17, 860-869
- 10 82. Fan, M., *et al.* (2004) Suppression of mitochondrial respiration through
11 recruitment of p160 myb binding protein to PGC-1alpha: modulation by p38
12 MAPK. *Genes Dev* 18, 278-289
- 13 83. Puigserver, P., *et al.* (1999) Activation of PPARgamma coactivator-1
14 through transcription factor docking. *Science* 286, 1368-1371
- 15 84. Ge, K., *et al.* (2002) Transcription coactivator TRAP220 is required for
16 PPAR gamma 2-stimulated adipogenesis. *Nature* 417, 563-567
- 17 85. Li, S., *et al.* (2008) Genome-wide coactivation analysis of PGC-1alpha
18 identifies BAF60a as a regulator of hepatic lipid metabolism. *Cell Metab* 8, 105-
19 117
- 20 86. Benton, C.R., *et al.* (2008) Modest PGC-1alpha overexpression in muscle
21 in vivo is sufficient to increase insulin sensitivity and palmitate oxidation in
22 subsarcolemmal, not intermyofibrillar, mitochondria. *J Biol Chem* 283, 4228-4240

- 1 87. Miura, S., *et al.* (2006) Overexpression of peroxisome proliferator-
- 2 activated receptor gamma co-activator-1alpha leads to muscle atrophy with
- 3 depletion of ATP. *Am J Pathol* 169, 1129-1139

4

Figure Legends

Fig. 1. Therapeutic potential of PGC-1 α . For a number of tissues, different functions of PGC-1 α have been described. A modulation of PGC-1 α expression and activity could have therapeutic implications in the diseases listed for each of these organs. Abbreviations: BAT, brown adipose tissue; CVD, cardiovascular diseases.

Fig. 2. Posttranslational modifications alter the specificity of PGC-1 α . The specificity of PGC-1 α to interact with its binding partners is altered by posttranslational modifications of the PGC-1 α protein. For example, phosphorylation of PGC-1 α by the AMP-dependent kinase (AMPK) primarily affects PGC-1 α target genes that are involved in cellular metabolism. Selective activation of neuromuscular junction (NMJ) genes in synaptic nuclei of the muscle fiber is achieved by different phosphorylations of PGC-1 α through motor neuron-derived signals. Abbreviations: OXPHOS, oxidative phosphorylation.

Fig. 3. Clinical modulation of PGC-1 α has to occur in a therapeutically beneficial window. Super- and subphysiological expression of PGC-1 α cause detrimental effects as described for heart and skeletal muscle. A pharmacological modulation should thus aim at a moderate increase or a normalization of pathologically dysregulated PGC-1 α . The therapeutic window might differ between tissues and contexts.

Fig. 1




Fig. 2




Fig. 3

